CLAIMS

1. Compounds of the general formula (I)

$$(R^{1})_{n1}$$
 $(R^{2})_{n2}$

in which:

R¹, which may be identical or different, are chosen independently from -Hal, -O-Alk, -N(Alk)₂, -NH-C(=O)-Alk, -O-C-Hal₃, -NO₂, -NH₂, -NHAlk, -C(=O)Hal, -C(=O)OAlk, -OH, -C(=O)-NH₂,-C(=O)-NHAlk, -C(=O)-N(Alk)₂, -NH-(C=O)-OAlk; -H, -CN, -Alk, -C(=O)Alk, -NAlk-C(O)-O-Alk and -NAlk₂;

n1 = integer between 1 and 5;

10 R², which may be identical or different, are chosen independently from -X-(C=O)-Y-(A)n and -CN;

n2 = integer between 1 and 5;

n = 0 when Y = Hal; n = 1 when Y = O or n = 2 when Y = N;

X = bond or -Alk-;

15 $Y = -O_{-}, -N_{-}$ or Hal;

25

A = -H, -Alk, -Alk-Ar, -Alk-Het, -Ar or -Het, Ar and Het optionally being substituted by Het or Hal;

 $R^3 = -H_1 - Alk_1 - NAlk_2 - NHAlk or - NH_2$

with the exception of the compounds for which:

• R^2 = -CN in position 4 (4-CN), n^2 = 1,

 $- R^3 = H$, n1 = 2 and $R^1 = (2-NHMe, 5-NO_2)$; or

 $- R^3 = Me, n1 = 1 \text{ and } R^1 = -NH-(C=O)-O-Me;$

• R^2 = 4-(-COOH), n2 = 1, and

- R^3 = (CH₂)₁₅-Me, n1 = 4, and R^1 = (2-OH,3,5-diCl,4-Et)

 $- R^3 = H$, n1 = 4, and $R^1 = (2-OH,3,5-diCl,4-Me)$

• $R^2 = 4$ -(-COOMe), $n^2 = 1$, $n^1 = 1$ and

 $-R^3 = H, R^1 = 2-(-C(=0)-CH_3)$

and also the addition salts thereof.

10

- 2. Compounds of the general formula (I) according to Claim 1, such that: $R^1 = -Hal$, -O-Alk, $-N(Alk)_2$, -NH-C(=O)-Alk, $-O-C-Hal_3$ or $-NO_2$.
- 5 3. Compounds according to Claim 1 or 2, in which n1 = 1.
 - 4. Compounds according to any one of the preceding claims, in which:

 R² = -(C=O)-NH₂, -COOH, -COHal, -(C=O)-OAlk, -CN, -(C=O)-NH-Het,

 -(C=O)-NH-Alk-Het, -Alk-(C=O)-NH-Phe, -Alk-(C=O)-NH-Het,

 -Alk-(C=O)-NH-Phe-Hal, -Alk-(C=O)-NH-Phe-Het.
 - 5. Compounds according to any one of the preceding claims, in which n2 = 1.
- 6. Compounds according to any one of the preceding claims, in which $R^3 = -H$ or -Alk.
 - 7. Compounds according to any one of the preceding claims in which $R^1 = -F$, -CI, -O-Me, -NMe₂, -NH-C(=O)-Me, -O-CF₃ or -NO₂.
- 20 8. Compounds according to any one of the preceding claims, in which: $R^2 = -CN, -COOH, -COCI, -(C=O)-OMe, -(C=O)-OEt, -(C=O)-NH_2, -(C=O)-NH-Py, -CH_2-(C=O)-NH-Py, -(C=O)-NH-(CH_2)_3-Im, -CH_2-(C=O)-NH-Phe, -CH_2-(C=O)-NH-Phe-F or -CH_2-(C=O)-NH-Phe-Morph.$
- 9. Compounds of the general formula (I) according to any one of the preceding claims, chosen from:
 - 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}-N-pyridin-3-ylbenzamide methyl 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}benzoate
 - 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}benzoic acid
- 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}benzoyl chloride 2-(4-{2-[(4-fluorophenyl)amino]-2-oxoethyl}phenoxy}-N-2-methoxyphenyl)acetamide

2-(4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}phenyl}-N-4-morpholin-4-ylphenyl)acetamide
2-[4-(2-anilino-2-oxoethyl)phenoxy]-N-(4-methoxyphenyl)acetamide
2-(4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}phenyl)-N-pyridin-3-ylethanamide
5 ethyl 4-{2-[(4-methoxyphenylamino]-2-oxoethoxy}benzoate
N-[3-(1h-imidazol-1-yl)propyl]-4-{2-[(methoxyphenyl)amino]-2-oxoethoxy}benzoate
methyl 4-{2-[(4-dimethylaminophenyl)amino]-2-oxoethoxy}benzoate

methyl 4-{2-[(4-N-acetylaminophenyl)amino]-2-oxoethoxy}benzoate 4-(2-oxo-2-{[4-(trifluoromethoxy)phenyl]amino}ethoxy)benzamide

2-(4-cyanophenoxy-N-[4-(trifluoromethoxy)phenyl]acetamide
4-{2-[(4-fluorophenyl)amino]-2-oxoethoxy}benzamide

4-{2-[(4-Nitrophenyl)amino]-2-oxoethoxy}(benzamide methyl 4-{2-[(3-chlorophenyl)amino]-2-oxoethoxy}benzoate methyl 4-{2-[(4-fluorophenyl)amino]-2-oxoethoxy}benzoate

methyl 4-(1-{[(4-fluorophenyl)amino]carbonyl}propoxy)benzoate and also the addition salts thereof.

10. Process for the preparation of the compounds of the general formula

$$(R^1)_{n1}$$
 $(R^2)_{n2}$

20

characterized in that a Smiles rearrangement is performed, starting with the compounds of the general formula (I)

$$(R^1)_{n1}$$
 $(R^2)_{n2}$

in which, in the general formulae (I) and (II):

 R^1 , which may be identical or different, are chosen independently from -Hal, -O-Alk, -N(Alk)₂, -NH-C(=O)-Alk, -O-C-Hal₃, -NO₂, -NH2, -NHAlk, -COOH, -C(=O)Hal, -C(=O)OAlk, -OH, -C(=O)-NH₂,-C(=O)-NHAlk, -C(=O)-N(Alk)₂, -N-(C=O)-OAlk; -H, -CN, -Alk, -C(=O)Alk, -NAlk-CO-OAlk and -NAlk₂,

5 n1 = integer between 1 and 5;

R², which may be identical or different, are chosen independently from-X-(C=O)-Y-(A)n and -CN;

n2 = integer between 1 and 5;

n = 0 when Y = Hal; n = 1 when Y = O or n = 2 when Y = N;

10 X = bond or -Alk-;

 $Y = -O_{-}, -N_{-}$ or Hal;

A = -H, -Alk, -Alk-Ar, -Alk-Het, -Ar or -Het, Ar and Het optionally being substituted by Het or Hal;

 $R^3 = -H$, -Aik, $-NAik_2$, -NHAik or $-NH_2$.

15

11. Process according to Claim 10, for which, in the general formulae (I) and (II),

 $R^1 = -Hal. -O-Alk. -N(Alk)_2, -NH-C(=O)-Alk. -O-C-Hal_3 or -NO_2;$

n1 = 1:

 $R^2 = -(C=O)-NH_2$, -COOH, -COHal, -(C=O)-OAlk, -CN, -(C=O)-NH-Het,

0 -(C=O)-NH-Alk-Het, -Alk-(C=O)-NH-Phe, -Alk-(C=O)-NH-Het,

-Alk-(C=O)-NH-Phe-Hal or -Alk-(C=O)-NH-Phe-Het,

n2 = 1:

 $R^3 = -H$ or -Alk.

25 12. Process according to Claim 10 or 11, for which, in the general formulae (I) and (II),

 R^1 = -F, -Cl, -O-Me, -NMe₂, -NH-C(=O)-Me, -O-CF₃ or -NO₂,

n1 = 1:

 $R^2 = -CN$, -COOH, -COCI, -(C=O)-OMe, -(C=O)-OEt, -(C=O)-NH₂, -(C=O)-NH-Py,

-CH₂-(C=O)-NH-Py, -(C=O)-NH-(CH₂)₃-Im, -CH₂-(C=O)-NH-Phe,

-CH₂-(C=O)-NH-Phe-F or -CH₂-(C=O)-NH-Phe-Morph,

n2 = 1;

 $R^3 = -H$ or Et

- 13. Process according to any one of Claims 10 to 12, for which the compounds of the general formula (I) are chosen from:
- 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}-N-pyridin-3-ylbenzamide methyl 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}benzoate
 - 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}benzoic acid
 - 4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}benzoyl chloride
 - 2-(4-{2-[(4-fluorophenyl)amino]-2-oxoethyl}phenoxy}-N-2-methoxyphenyl)acetamide
- 2-(4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}phenyl}-N-4-morpholin-4-ylphenyl)-acetamide
 - 2-[4-(2-anilino-2-oxoethyl)phenoxy]-N-(4-methoxyphenyl)acetamide
 - 2-(4-{2-[(4-methoxyphenyl)amino]-2-oxoethoxy}phenyl)-N-pyridin-3-ylethanamide ethyl 4-{2-[(4-methoxyphenylamino]-2-oxoethoxy}benzoate
- N-[3-(1h-imidazol-1-yl)propyl]-4-{2-[(methoxyphenyl)amino]-2-oxoethoxy}benzamide methyl 4-{2-[(4-dimethylaminophenyl)amino]-2-oxoethoxy}benzoate methyl 4-{2-[(4-N-acetylaminophenyl)amino]-2-oxoethoxy}benzoate 4-(2-oxo-2-{[4-(trifluoromethoxy)phenyl]amino}ethoxy)benzamide
 - 2-(4-cyanophenoxy-N-[4-(trifluoromethoxy)phenyl]acetamide
- 20 4-{2-[(4-fluorophenyl)amino]-2-oxoethoxy}benzamide
 - 4-{2-[(4-Nitrophenyl)amino]-2-oxoethoxy}(benzamide
 - methyl 4-{2-[(3-chlorophenyl)amino]-2-oxoethoxy}benzoate
 - methyl 4-{2-[(4-fluorophenyl)amino]-2-oxoethoxy}benzoate
 - methyl 4-(1-{[(4-fluorophenyl)amino]carbonyl}propoxy)benzoate
- 25 and also the addition salts thereof.
 - 14. Process according to any one of Claims 10 to 13, which is performed in basic medium, in a suitable solvent, with stirring and at reflux.
- 15. Process according to Claim 14, for which the base is chosen from sodium hydroxide, potassium hydroxide and potassium carbonate.

- 16. Process according to Claim 14 or 15, for which the solvent is chosen from amides.
- 5 17. Process according to any one of Claims 14 to 16, which is performed at between 70 and 160°C.
 - 18. Process according to any one of Claims 5 to 12, for which the compound of the general formula (II) is obtained by using the compounds of the general formulae (III) and (IV), as defined according to any one of Claims 21 to 24, without isolation of the intermediate compound (I) formed.
 - 19. Use of the compounds of the general formula (I)

$$(R^{1})_{n1}$$
 $(R^{2})_{n2}$

for the preparation of compounds of the general formula (II):

20

$$(R^{1})_{n1} \xrightarrow{\qquad \qquad } (R^{2})_{n2}$$

in which R¹, R², R³, n1 and n2 are as defined in Claims 10 to 12.

20. Process for the preparation of the compounds of the general formula (I)

$$(R^{1})_{n1}$$
 $(R^{2})_{n2}$

characterized in that the compounds of the general formulae (III) and (IV) are used:

$$(\mathbb{R}^1)_{n1} \xrightarrow{\qquad \qquad } \mathbb{R}^3$$

in which R¹, R², R³, n1 and n2 are as defined in Claims 10 to 12, and for which the process is performed in basic medium, in a suitable solvent, with stirring and at reflux.

- 21. Process according to Claim 20, for which the base is chosen from sodium hydroxide, potassium hydroxide and potassium carbonate.
- 10 22. Process according to Claim 21, for which the solvent is chosen from amides.
 - 23. Process according to either of Claims 21 and 22, which is performed at a temperature of between room temperature and 130°C.
- 15 24. Process according to any one of Claims 21 to 23, for which the compound of the general formula (III) is obtained by using the compounds of the general formulae (V) and (VI):

$$(R^{1})_{n1} \xrightarrow{\text{NH}_{2}} CI \xrightarrow{\text{CI}} CI$$

$$(V) \qquad (VI)$$

20 25. Compounds of the general formula (I)

$$(R^{1})_{n1}$$
 $(R^{2})_{n2}$

in which R1, R2, R3, n1 and n2 are as defined in Claims 10 to 12,

WO 2005/051892 PCT/EP2004/012541 28

which can be obtained via the process according to any one of Claims 21 to 24.